
















Product Name / Pest	Crops Affected	No. of Traps	Field Viability	Ideal Time to Install Trap	Recommended Traps
Melon Fly Lure (<i>Bactrocera cucurbitae</i>)	 Cucumber, Bottle Gourd, Ridge Gourd, Bitter Gourd, Snake Gourd, Watermelon, Muskmelon, Pumpkins, etc.	6	60 Days	30 days after sowing	 Insect Fly Trap
Fruit Fly Lure (<i>Bactrocera dorsalis</i>)	 Mango, Guava, Lemon, etc.	6	60 Days	10 days after flowering	
Rhinoceros Beetle Lure (<i>Oryctes rhinoceros</i>)	 Coconut, Date Palms, Oil Palms, Areca Nut	2	120 Days	Anytime of the year	 Insect Bucket Trap
Red Palm Weevil Lure (<i>Rhynchophorus ferrugineus</i>)	 Coconut, Date Palms, Oil Palms, Areca Nut	2	180 Days	Anytime of the year	
Pink Bollworm Lure (<i>Pectinophora gossypiella</i>)	 Cotton	12	60 Days	30 days after sowing	 Insect Funnel Trap
Fall Armyworm Lure (<i>Spodoptera frugiperada</i>)	 Corn, Cotton, Rice, Millets, etc.	16	30 Days	10 days after sowing	
American Bollworm Lure (<i>Helicoverpa armigera</i>)	 Cotton, Pigeon Pea, Chickpea, Tomato, Capsicum, Corn, Green Pea, Grapes, etc.	12	60 Days	Chickpea At time of sowing Pigeon pea 60 days after sowing	
Tobacco Caterpillar Lure (<i>Spodoptera litura</i>)	 Cabbage, Tobacco, Soybean, Cotton, Grapes, Chillies, Sunflower, Groundnut, etc.	12	60 Days	After transplanting	
Rice Yellow Stem Borer Lure (<i>Scirpophaga incertulas</i>)	 Rice	16	60 Days	After transplanting	
Brinjal Fruit & Shoot Borer Lure (<i>Leucinodes orbonalis</i>)	 Brinjal	16	60 Days	After transplanting	
Diamondback Moth Lure (<i>Plutella xylostella</i>)	 Cabbage, Cauliflower	30	60 Days	After transplanting	 Insect Water Trap
Tomato Leafminer Lure (<i>Tuta absoluta</i>)	 Tomato	16	60 Days	After transplanting	